



References

NICU: CLABSI GUIDELINES
PAGE 7 of 8 | [ALL PAGES](#) ↓

References from the Recommendations for Prevention and Control of Infections in NICU Patients: CLABSI

AT A GLANCE

Review references from the Recommendations for Prevention and Control of Infections in NICU Patients: CLABSI guideline.

ON THIS PAGE

References

References

1. The Healthcare Infection Control Practices Advisory Committee (HICPAC). Core Infection Prevention and Control Practices for Safe Healthcare Delivery in All Settings – Recommendations of the Healthcare Infection Control Practices Advisory Committee. The Centers for Disease Control and Prevention. Updated December 27, 2018. Accessed January 20, 2020. <https://www.cdc.gov/hicpac/recommendations/core-practices.html>
2. Centers for Disease Control and Prevention. Guidelines for the Prevention of Intravascular Catheter-Related Infections. Centers for Disease Control and Prevention,. Updated November 5, 2015. Accessed August 14, 2020. <https://www.cdc.gov/infectioncontrol/guidelines/bsi/index.html>
3. Kaufman DA, Blackman A, Conaway MR, Sinkin RA. Nonsterile glove use in addition to hand hygiene to prevent late-onset infection in preterm infants: Randomized clinical trial. *JAMA Pediatrics*. 01 Oct 2014;168(10):909-916. doi:<http://dx.doi.org/10.1001/jamapediatrics.2014.953>
4. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. Jun 2008;36(5):309-32. doi:10.1016/j.ajic.2008.03.002
5. Arnts IJ, Schrijvers NM, van der Flier M, Groenewoud JM, Antonius T, Liem KD. Central line bloodstream infections can be reduced in newborn infants using the modified Seldinger technique and care bundles of preventative measures. Observational Study. *Acta Paediatr*. Apr 2015;104(4):e152-7. doi:<https://dx.doi.org/10.1111/apa.12915>
6. Bhandari V, Eisenfeld L, Lerer T, Holman M, Rowe J. Nosocomial sepsis in neonates with single lumen vascular catheters. *Indian Journal of Pediatrics*. July/August 1997;64(4):529-535.
7. Chien LY, Macnab Y, Aziz K, Andrews W, McMillan DD, Lee SK. Variations in central venous catheter-related infection risks among Canadian neonatal intensive care units. *Pediatric Infectious Disease Journal*. 2002;21(6):505-511. doi:<http://dx.doi.org/10.1097/00006454-200206000-00006>
8. de Brito CS, de Brito DV, Abdallah VO, Gontijo Filho PP. Occurrence of bloodstream infection with different types of central vascular catheter in critically neonates. *J Infect*. Feb 2010;60(2):128-32. doi:<https://dx.doi.org/10.1016/j.jinf.2009.11.007>
9. Geldenhuys CD, A., Jenkins A, Bekker A, Jenkins A, Bekker A. Central-line-associated bloodstream infections in a resource-limited South African neonatal intensive care unit. *South African Medical Journal*. 2017;107(9):758-762. doi:<http://dx.doi.org/10.7196/SAMJ.2017.v107i9.12124>
10. Greenberg RG, Cochran KM, Smith PB, et al. Effect of Catheter Dwell Time on Risk of Central Line-Associated Bloodstream Infection in Infants. *Pediatrics*. 2015;136(6):1080-1086. doi:10.1542/peds.2015-0573

11. Sanderson E, Yeo KT, Wang AY, et al. Dwell time and risk of central-line-associated bloodstream infection in neonates. *Journal of Hospital Infection*. November 2017;97(3):267-274. doi:<http://dx.doi.org/10.1016/j.jhin.2017.06.023> 
12. Shalabi MA, Yoon M, Aziz E, Shoo K, Shah L, Prakesh S. Risk of Infection Using Peripherally Inserted Central and Umbilical Catheters in Preterm Neonates. *Pediatrics*. 2015;136(6):1073-1079. doi:10.1542/peds.2015-2710
13. Soares BN, Pissarra S, Rouxinol-Dias AL, Costa S, Guimaraes H. Complications of central lines in neonates admitted to a level III Neonatal Intensive Care Unit. In Press. *Journal of Maternal-Fetal and Neonatal Medicine*. 26 Jul 2017:1-7. doi:<http://dx.doi.org/10.1080/14767058.2017.1355902> 
14. Chenoweth KB, Guo JW, Chan B. The Extended Dwell Peripheral Intravenous Catheter Is an Alternative Method of NICU Intravenous Access. *Adv Neonat Care*. 2018;18(4):295-301.
15. Konstantinidi A, Sokou R, Panagiotounakou P, et al. Umbilical Venous Catheters and Peripherally Inserted Central Catheters: Are They Equally Safe in VLBW Infants? A Non-Randomized Single Center Study. *Medicina*. 2019;55(8):06.
16. Bashir RA, Swarnam K, Vayalthrikkovil S, Yee W, Soraisham AS. Association between Peripherally Inserted Central Venous Catheter Insertion Site and Complication Rates in Preterm Infants. *American Journal of Perinatology*. 01 Aug 2016;33(10):945-950. doi:<http://dx.doi.org/10.1055/s-0036-1582127> 
17. Breschan C, Platzer M, Jost R, Schaumberger F, Stettner H, Likar R. Comparison of catheter-related infection and tip colonization between internal jugular and subclavian central venous catheters in surgical neonates. *Anesthesiology*. December 2007;107(6):946-953. doi:<http://dx.doi.org/10.1097/01.anes.0000291443.78166.98> 
18. Hoang V, Sills J, Chandler M, Busalani E, Clifton-Koeppel R, Modanlou HD. Percutaneously inserted central catheter for total parenteral nutrition in neonates: complications rates related to upper versus lower extremity insertion. *Pediatrics*. 2008;121(5):e1152-9.
19. Tsai MH, Chu SM, Lien R, et al. Complications associated with 2 different types of percutaneously inserted central venous catheters in very low birth weight infants. *Infect Control Hosp Epidemiol*. Mar 2011;32(3):258-66. doi:<https://dx.doi.org/10.1086/658335> 
20. Tsai MH, Lien R, Wang JW, et al. Complication rates with central venous catheters inserted at femoral and non-femoral sites in very low birth weight infants. *Pediatric Infectious Disease Journal*. Nov 2009;28(11):966-70. doi:<https://dx.doi.org/10.1097/INF.0b013e3181aa3a29> 
21. Vegunta RK, Loethen P, Wallace LJ, Albert VL, Pearl RH. Differences in the outcome of surgically placed long-term central venous catheters in neonates: Neck vs groin placement. *Journal of Pediatric Surgery*. January 2005;40(1):47-51. doi:<http://dx.doi.org/10.1016/j.jpedsurg.2004.09.015> 
22. Wrightson DD. Peripherally inserted central catheter complications in neonates with upper versus lower extremity insertion sites. Comparative Study Evaluation Studies. *Adv Neonat Care*. Jun 2013;13(3):198-204. doi:<https://dx.doi.org/10.1097/ANC.0b013e31827e1d01> 
23. Elmekawi A, Maulidi H, Mak W, Aziz A, Lee KS. Outcomes of upper extremity versus lower extremity placed peripherally inserted central catheters in a medical-surgical neonatal intensive care unit. *Journal of Neonatal-Perinatal Medicine*. 2019;12(1):57-63.
24. Garcia H, Romano-Carro B, Miranda-Novales G, Gonzalez-Cabello HJ, Nunez-Enriquez JC. Risk Factors for Central Line-Associated Bloodstream Infection in Critically Ill Neonates. *Indian Journal of Pediatrics*. 2019;86(4):340-346.
25. Litz CN, Tropf JG, Danielson PD, Chandler NM. The idle central venous catheter in the NICU: When should it be removed? *Journal of Pediatric Surgery*. 2018;53(7):1414-1416.
26. Khilnani P, Goldstein B, Todres ID. Double lumen umbilical venous catheters in critically ill neonates: a randomized prospective study. *Critical care medicine*. Nov 1991;19(11):1348-51. doi:10.1097/00003246-199111000-00007
27. Levit OL, Shabanova V, Bizzarro MJ. Umbilical catheter-associated complications in a level IV neonatal intensive care unit. *J Perinatol*. Apr 2020;40(4):573-580. doi:10.1038/s41372-019-0579-3
28. Ratz D, Hofer T, Flanders SA, Saint S, Chopra V. Limiting the Number of Lumens in Peripherally Inserted Central Catheters to Improve Outcomes and Reduce Cost: A Simulation Study. *Infect Control Hosp Epidemiol*. Jul 2016;37(7):811-7. doi:10.1017/ice.2016.55
29. Garland JS, Alex CP, Uhing MR, Peterside IE, Rentz A, Harris MC. Pilot trial to compare tolerance of chlorhexidine gluconate to povidone-iodine antiseptics for central venous catheter placement in neonates. *J Perinatol*. Dec 2009;29(12):808-13. doi:10.1038/jp.2009.161
30. Sankar MJ, Paul VK, Kapil A, et al. Does skin cleansing with chlorhexidine affect skin condition, temperature and colonization in hospitalized preterm low birth weight infants?: a randomized clinical trial. *J Perinatol*. Dec 2009;29(12):795-801. doi:10.1038/jp.2009.110

31. Cleves D, Pino J, Patino JA, Rosso F, Velez JD, Perez P. Effect of chlorhexidine baths on central-line-associated bloodstream infections in a neonatal intensive care unit in a developing country. *J Hosp Infect*. Nov 2018;100(3):e196-e199. doi:10.1016/j.jhin.2018.03.022
32. Westling T, Cowden C, Mwananyanda L, et al. Impact of chlorhexidine baths on suspected sepsis and bloodstream infections in hospitalized neonates in Zambia. *International Journal of Infectious Diseases*. 2020;96:54-60. doi:<https://dx.doi.org/10.1016/j.ijid.2020.03.043> 
33. Quach C, Milstone AM, Perpete C, Bonenfant M, Moore DL, Perreault T. Chlorhexidine bathing in a tertiary care neonatal intensive care unit: impact on central line-associated bloodstream infections. *Infect Control Hosp Epidemiol*. Feb 2014;35(2):158-63. doi:10.1086/674862
34. Mahieu LM, De Dooy JJ, Lenaerts AE, Ieven MM, De Muynck AO. Catheter manipulations and the risk of catheter-associated bloodstream infection in neonatal intensive care unit patients. *J Hosp Infect*. May 2001;48(1):20-6. doi:10.1053/jhin.2000.0930
35. Filippi L, Pezzati M, Di Amario S, Poggi C, Pecile P. Fusidic acid and heparin lock solution for the prevention of catheter-related bloodstream infections in critically ill neonates: a retrospective study and a prospective, randomized trial. *Pediatr Crit Care Med*. Nov 2007;8(6):556-62. doi:10.1097/01.PCC.0000288711.46009.58
36. Garland JS, Alex CP, Henrickson KJ, McAuliffe TL, Maki DG. A vancomycin-heparin lock solution for prevention of nosocomial bloodstream infection in critically ill neonates with peripherally inserted central venous catheters: a prospective, randomized trial. *Pediatrics*. Aug 2005;116(2):e198-205. doi:10.1542/peds.2004-2674.
37. Seliem W. Amikacin-heparin lock for prevention of catheter-related bloodstream infection in neonates with extended umbilical venous catheters use: A randomized controlled trial. *Journal of neonatal-perinatal medicine*. 2010;3(1):33-41.
38. Butler-O'Hara M, Buzzard CJ, Reubens L, McDermott MP, DiGrazio W, D'Angio CT. A randomized trial comparing long-term and short-term use of umbilical venous catheters in premature infants with birth weights of less than 1251 grams. *Pediatrics*. Jul 2006;118(1):e25-35. doi:10.1542/peds.2005-1880.
39. Butler-O'Hara M, D'Angio CT, Hoey H, Stevens TP. An evidence-based catheter bundle alters central venous catheter strategy in newborn infants. *J Pediatr*. Jun 2012;160(6):972-7.e2. doi:10.1016/j.jpeds.2011.12.004.
40. Vachharajani AJ, Vachharajani NA, Morris H, et al. Reducing peripherally inserted central catheters in the neonatal intensive care unit. *Journal of Perinatology*. 01 Apr 2017;37(4):409-413. doi:<http://dx.doi.org/10.1038/jp.2016.243> 
41. Hsu JF, Tsai MH, Huang HR, Lien R, Chu SM, Huang CB. Risk factors of catheter-related bloodstream infection with percutaneously inserted central venous catheters in very low birth weight infants: A center's experience in Taiwan. *Pediatrics and Neonatology*. December 2010;51(6):336-342. doi:<http://dx.doi.org/10.1016/S1875-9572%2810%2960065-4> 
42. Milstone AM, Reich NG, Advani S, et al. Catheter dwell time and CLABSI in neonates with PICCs: a multicenter cohort study. Evaluation Studies Multicenter Study Research Support, N.I.H., Extramural. *Pediatrics*. Dec 2013;132(6):e1609-15. doi:<https://dx.doi.org/10.1542/peds.2013-1645> 
43. Njere I, Islam S, Parish D, Kuna J, Keshtgar AS. Outcome of peripherally inserted central venous catheters in surgical and medical neonates. Conference Paper. *Journal of Pediatric Surgery*. May 2011;46(5):946-950. doi:<http://dx.doi.org/10.1016/j.jpedsurg.2011.02.037> 
44. Ohki Y, Maruyama K, Harigaya A, Kohno M, Arakawa H. Complications of peripherally inserted central venous catheter in Japanese neonatal intensive care units. Comparative Study Multicenter Study. *Pediatr Int*. Apr 2013;55(2):185-9. doi:<https://dx.doi.org/10.1111/ped.12033> 
45. Rangel UV, Gomes Junior SC, Costa AM, Moreira MEL. Variables associated with peripherally inserted central catheter related infection in high risk newborn infants. *Rev Lat Am Enfermagem*. Oct 2014;22(5):842-7.
46. Sengupta A, Lehmann C, Diener-West M, Perl TM, Milstone AM. Catheter duration and risk of CLA-BSI in neonates with PICCs. Comparative Study Research Support, N.I.H., Extramural. *Pediatrics*. Apr 2010;125(4):648-53. doi:<https://dx.doi.org/10.1542/peds.2009-2559> 
47. Taylor T, Massaro A, Williams L, et al. Effect of a dedicated percutaneously inserted central catheter team on neonatal catheter-related bloodstream infection. *Adv Neonatal Care*. Apr 2011;11(2):122-8. doi:10.1097/ANC.0b013e318210d059
48. Holzmann-Pazgal G, Kubanda A, Davis K, Khan AM, Brumley K, Denson SE. Utilizing a line maintenance team to reduce central-line-associated bloodstream infections in a neonatal intensive care unit. *Journal of Perinatology*. Apr 2012;32(4):281-6. doi:<https://dx.doi.org/10.1038/jp.2011.91> 
49. Centers for Disease Control and Prevention. Checklist for Prevention of Central Line-associated Blood Stream Infection. Accessed December 1, 2020. <https://www.cdc.gov/hai/pdfs/bsi/checklist-for-clabsi.pdf> 

50. Fisher D, Cochran KM, Provost LPP, J. , et al. Reducing central line-associated bloodstream infections in North Carolina NICUs. Clinical Trial Multicenter Study Research Support, Non-U.S. Gov't. *Pediatrics*. Dec 2013;132(6):e1664-71. doi:<https://dx.doi.org/10.1542/peds.2013-2000> 
51. Savage T, Hodge DE, Pickard K, Myers P, Powell K, Cayce JM. Sustained Reduction and Prevention of Neonatal and Pediatric Central Line-Associated Bloodstream Infection Following a Nurse-Driven Quality Improvement Initiative in a Pediatric Facility. *JAVA – Journal of the Association for Vascular Access*. 2018;23(1):30-41.
52. Balla KC, Rao SPN, Arul C, et al. Decreasing Central Line-associated Bloodstream Infections Through Quality Improvement Initiative. *Indian Pediatrics*. 2018;55(9):753-756.
53. Harms K, Herting E, Kron M, Schiffmann H, Schulz-Ehlbeck H. Randomized, controlled trial of amoxicillin prophylaxis for prevention of catheter-related infections in newborn infants with central venous silicone elastomer catheters. *J Pediatr*. Oct 1995;127(4):615-9. doi:10.1016/s0022-3476(95)70126-5
54. Elhassan NO, Stevens TP, Gigliotti F, Hardy DJ, Cole CA, Sinkin RA. Vancomycin usage in central venous catheters in a neonatal intensive care unit. *Pediatr Infect Dis J*. Mar 2004;23(3):201-6.
55. Ocete E, Ruiz-Extremuera A, Goicoechea A, et al. Low-dosage prophylactic vancomycin in central-venous catheters for neonates. *Early Hum Dev*. Dec 1998;53 Suppl:S181-6.
56. Spafford PS, Sinkin RA, Cox C, Reubens L, Powell KR. Prevention of central venous catheter-related coagulase-negative staphylococcal sepsis in neonates. *J Pediatr*. Aug 1994;125(2):259-63. doi:10.1016/s0022-3476(94)70208-x
57. Birch P, Ogden S, Hewson M. A randomised, controlled trial of heparin in total parenteral nutrition to prevent sepsis associated with neonatal long lines: the Heparin in Long Line Total Parenteral Nutrition (HILLTOP) trial. Randomized Controlled Trial Research Support, Non-U.S. Gov't. *Arch Dis Child Fetal Neonatal Ed*. Jul 2010;95(4):F252-7. doi:<https://dx.doi.org/10.1136/adc.2009.167403> 
58. Kamala F, Boo NY, Cheah FC, Birinder K. Randomized controlled trial of heparin for prevention of blockage of peripherally inserted central catheters in neonates. *Acta Paediatrica, International Journal of Paediatrics*. 2002;91(12):1350-1356.
59. Shah PS, Kalyn A, Satodia P, et al. A randomized, controlled trial of heparin versus placebo infusion to prolong the usability of peripherally placed percutaneous central venous catheters (PCVCs) in neonates: the HIP (Heparin Infusion for PCVC) Study. *Pediatrics*. 2007;119(1):e284-91.
60. Uslu S, Ozdemir H, Comert S, Bolat F, Nuhoglu A. The effect of low-dose heparin on maintaining peripherally inserted percutaneous central venous catheters in neonates. *J Perinatol*. Dec 2010;30(12):794-9. doi:10.1038/jp.2010.46
61. Payne N, Barry J, Berg W, et al. Sustained Reduction in Neonatal Nosocomial Infections Through Quality Improvement Efforts. *Pediatrics*. 2012;129(1):e1-e9.
62. Stoll BJ, Hansen NI, Adams-Chapman I, et al. Neurodevelopmental and growth impairment among extremely low-birth-weight infants with neonatal infection. *JAMA*. Nov 17 2004;292(19):2357-65. doi:10.1001/jama.292.19.2357
63. Schulman J, Stricof R, Stevens TP, et al. Statewide NICU central-line-associated bloodstream infection rates decline after bundles and checklists. *Pediatrics*. Mar 2011;127(3):436-44. doi:10.1542/peds.2010-2873
64. Centers for Disease Control and Prevention. The Targeted Assessment for Prevention (TAP) Strategy. Updated June 21, 2019. Accessed August 15, 2020. <https://www.cdc.gov/hai/prevent/tap.html>
65. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. Apr 2011;64(4):383-94. doi:10.1016/j.jclinepi.2010.04.026
66. Schunemann HJ, Oxman AD, Brozek J, et al. GRADE: assessing the quality of evidence for diagnostic recommendations. *ACP J Club*. Dec 16 2008;149(6):2.
67. The Healthcare Infection Control Practices Advisory Committee (HICPAC). Update to the CDC and the HICPAC Recommendation Categorization Scheme for Infection Control and Prevention Guideline Recommendations Updated October 8, 2019. Accessed August 14, 2020. <https://www.cdc.gov/hicpac/workgroup/recommendation-scheme-update.html>
68. Arnts IJ, Groenewoud LM, Liem JMM, Djien K. Comparison of Complication Rates Between Umbilical and Peripherally Inserted Central Venous Catheters in Newborns. *JOGNN: Journal of Obstetric, Gynecologic & Neonatal Nursing*. 2014;43(2):205-215. doi:10.1111/1552-6909.12278

69. Taylor JE, McDonald SJ, Earnest A, et al. A quality improvement initiative to reduce central line infection in neonates using checklists. *Eur J Pediatr*. May 2017;176(5):639-646. doi:<https://dx.doi.org/10.1007/s00431-017-2888-x> 

70. Esaiassen E, Fjalstad JW, Juvet LK, van den Anker JN, Klingenberg C. Antibiotic exposure in neonates and early adverse outcomes: a systematic review and meta-analysis. *J Antimicrob Chemother*. Jul 1 2017;72(7):1858-1870. doi:10.1093/jac/dkx088

71. Cantey JB, Huffman LW, Subramanian A, et al. Antibiotic Exposure and Risk for Death or Bronchopulmonary Dysplasia in Very Low Birth Weight Infants. *J Pediatr*. Feb 2017;181:289-293 e1. doi:10.1016/j.jpeds.2016.11.002

READ NEXT
Acronyms and Abbreviations



TABLE OF CONTENTS
NICU: CLABSI GUIDELINES

1. Authors, Contributors and Acknowledgments	5. Methods
2. Executive Summary	6. Evidence Summaries
3. Summary of Recommendations	7. References
4. Introduction	8. Acronyms and Abbreviations

MARCH 19, 2024

 SOURCES

SHARE

CONTENT SOURCE:
[National Center for Emerging and Zoonotic Infectious Diseases \(NCEZID\)](#)

RELATED PAGES

- [Prevention and Control of Infections in Neonatal Intensive Care Unit Patients: Central Line-associated Blood Stream Infections](#)
- [Summary of Recommendations](#)
- [Introduction](#)
- [Methods](#)
- [Evidence Summaries](#)
- [Acronyms and Abbreviations](#)